

#### PAMANTASAN NG LUNGSOD NG MAYNILA

(University of the City of Manila) Intramuros, Manila

## **REQUEST FOR PROPOSAL**

Date:	
RFQ No.:	

Name of the Company:	
Address:	
Business Permit No.:	
TIN No.:	

Please quote your best offer for the item/s described below, subject to the Terms and Conditions provided at the back portion of this request for quotation. Submit your quotation duly signed by you or your duly authorized representative not later than \_\_\_\_

Open quotations may be submitted, manually or through facsimile or email at the address and contact number indicated below.

> SIGNED **ALBERT S. DELA CRUZ** Chief, Procurement Office

After having carefully read and accepted the Terms and Conditions, I/We submit our quotation/s for the item/s as follows:

				OFFER					
	Item Description	QTY	Approved Compliance witi Budget for Price Price Specifications (ABC) (please check)	Price		nnical ications	Remarks		
				QTY	Unit Price	Total Price	Yes	No	
	Software Authorized User License + 12- are Subscription & Support (S & S)	1Lot	150,000.00						
Background	PLM is in need of a Mathematics Software that is loaded with powerful analytic techniques and time-saving features to help PLM quickly and easily find new insights in institutional data that lead to more accurate predictions and achieve better outcomes for the University.								
Module Description	The Module should enable PLM to take a quick look at data, formulate hypotheses for additional testing, and then carry out statistical and analytical procedures to help clarify relationships between variables, create clusters, identify trends and make predictions.								
built-in techni Analyze data procedures Easily build capabilities Discover new mapping capa Quickly build customized d	Quickly access and accurately analyze massive data sets and manage the data for analysis through ques with comprehensive range of statistical charts with sophisticated reporting insights in the data with tables, graphs, bilities, cubes and pivoting technology dialog boxes or let advanced users create ialog boxes that make your organization's r and more efficient								
	Descriptive Statistics ensive software solution includes a wide cedures and test to solve business and enges.								
· Cross tabul marginal, te	ations: Counts, percentages, residuals, st of independence, tests of linear easure of linear association, ordinal data								

measure, normal by interval measure, measure of seven in the seven interval interva						
<pre>index tube: Contact, protectings, usid and contactors representations: Propundity: Contact, protectings, usid and contactors representations: Procession: Contactors of dispersion. Decentation: Contactors of d</pre>	measures, nominal by interval measure, measure of					
<pre>index turks. Frequencies: Const, performance, durations of the constructions of the theory. Specific durations and correct Decryption: Central tensory. Specific durations and correct Decryption: Central tensory. Specific durations and correct constructions for the specific duration of the specific duration of the specific durations and correct constructions for the specific duration of the specific duration of the specific duration of the specific duration of the specific duration of the specific duration of the speci</pre>	agreement, relative risk estimates for case control and					
i frequentità ci cuità precision, vidi ad cumulativa preprintaria ci cuità precision, distribution ad 2 sciene. L'accordinate distribution ad 2 sciene. Concordinate distribution ad 2 science. Concordinate distribut	•					
processings, cardial forshorm, dispersion, distribution and a processing status status; dispersion, dispersion, cardial of a writion, is becarging and a transfer of a writion, is cardial of a writion, is cardial of a writion, is cardial of a writion, is cardial of a writion, is cardial of a writion, is cardial of a writion, is cardial of a writion, is cardial of a writ						
and percention evides. Description: Control in the evides in the evides of the evides of the evides in the evide						
indexignore: Central tendency, disputsion, distribution additional control of the surf substrates and energies studies devices. Conference without the substrates and energies studies. Conference without the substrates and energies and substrates provide subscrates and the substrates provide conference without the subscrates and the substrate provide subscrates and the subscrates. Conference without the subscrates and the subscrates provide subscrates and the subscrates provide subscrates and the subscrates. The first subscrates and and subscrates. The first subscrates provide subscrates and the subscrates provide subscrates and the subscrates provide subscrates and the subscrates and the subscrates and and the subscrates and						
and zoors. Decryption proves that statistics: Confliction of dispersion, coverage frames. Unclose whether to use homemoin or generation council to large draw is the homemoin draw is the home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home draw is the heads of home draw is the home dra						
indexting and variables rules statistics: Confident of dispersion, compare order theory. Compared theory	· Descriptive: Central tendency, dispersion, distribution					
indificant variation, purce patient differential and provide a block definition. Second the means, testis function, compare tai independent second the means, testis function, compare tai independent second the means, testis function, compare tai independent test. Monte tai second tai means testis choices you model and the testis second tai testis second tai testis second definition of the land tai testis second testis second tai testis second tai testis second testis testis second tai testis second tai testis te	and Z scores.					
indificant variation, purce patient differential and provide a block definition. Second the means, testis function, compare tai independent second the means, testis function, compare tai independent second the means, testis function, compare tai independent test. Monte tai second tai means testis choices you model and the testis second tai testis second tai testis second definition of the land tai testis second testis second tai testis second tai testis second testis testis second tai testis second tai testis te	· Descriptive ratio statistics: Coefficient of dispersion.					
inverse backvice devinate Granter means: Obvice white it is use harmonic or parage trackticles, pureter simple studiets: or oversample test. - ALCVA and ALCCVA-Conduct contrast, range, and part is used on four types of the simple requires on the simple test of the simple studiets: or oversample test. - ALCVA and ALCCVA-Conduct contrast, range, and part is used on four types of the simple requires on the simple - ALCVA and ALCCVA-Conduct contrast, range, and part - ALCVA and range contrast contrast, range and part - ALCVA and range contrast contrast, range and part - ALCVA and ALCVA-Conduct contrast, range and part - ALCVA and range contrast contrasts within a - of alcer contrast, range and part - alcer contrast, range						
Compare means: Choose whether to use humonics or promotion means the laberity compare and underpredent terms. Another and another statistics or underpredent terms. Another and another statistics or underpredent terms. Statistics and means and terms of the statistics of the statistics of the terms. The another statistics and the statistics and terms and terms of the statistics of the statistics and the statistics of terms terms and terms of the statistics and the statistics of terms terms and terms of the statistics and the statistics of terms terms and terms of terms of the statistics and the statistics of terms terms and terms of ter						
jeonetic meanse test linearing: compare val independent units linearing and sample statistics, one one ample line. Out Sets hangle for fields and sample statistics one and model based on for types of the sam of account of the same factor measure; group description statistics, these you model based on for types of the sam of account of the same factor ensign of analytic covariance with up to 21 methods. Ground analytics of the same factor and the same factor statistics of the same factor and the same factor ensign of analytic covariance with up to 21 methods. Group and the same factor and the same factor statistics of the same factor statistics of t	5					
imple statistic, pandy: for distribution or one-imple test. Are test; analyze, fund offers, and andrem offers measure; group distribution statistic or unbalanced based on four type of the sum of searces proceeding: perform lack offersit, Chook balance of unbalanced consistence tests; Chook balance of unbalanced tests; Chook balance of performance, tests; Chook balance of performance, tes						
test AdVA and AdVCAD. Conduct contrast, engry, and post the mean opping involves tables, the two provided bases of Not Types of Heaution 4 subseq provided perform tack of Heaution 4 subseq provided subseq of Not Types of Heaution 4 subseq subseq of Heaution 4 subseq subseq of Heaution 4 subseq subseq of Not Types of Heaution 4 subseq subseq of Heaution 4 subseq subseq of Heaution 4 subseq subseq of A subseq Subseq of Not Types of Heaution 4 subseq subseq of A subseq Subseq of Not Subseq subseq of A subseq Subseq of Not Types of Heaution 4 subseq subseq of A subseq Subseq of Not Types of Heaution 4 subseq subseq of A subseq Subseq of Not Types of Heaution 4 subseq subseq of A subseq Subseq of Not Types of Heaution 4 subseq subseq of A subseq Subseq of Not Types of Heaution 4 subseq subseq of A subseq Subseq of Not Types of Heaution 4 subseq subseq of Heaution 4 subseq Not Types of Not Types of Heaution 4 subseq subseq of Heaution 4 subseq Not Types of Not Types of Heaution 4 subseq subseq of Heaution 4 subseq Not Types of Not Types of Heaution 4 subseq subseq Not Types of Not Types of Heaution 4 subseq subseq Not Types of Not Types of Heaution 4 subseq subseq	geometric means; test linearity; compare via independent					
-MOVA and ANCOVA Conduct contrast, regine, and pot hor: test analyse for defects and normality between meaning graph description stratustics, choose you model perform load of energy, how has also on unbalanced design, and wayter exventions with up to D methods. Concretion: Test of behaviors of mission many between meaning, the mission many or design many between meaning, the mission many or design many between meaning. The mission many or design many between meaning, the mission many or design many between meaning. The mission many or design many between meaning, the mission many or design many between meaning. The mission many or design many between meaning of the mission many of the mission many between meaning of the mission the test the mission many of the mission of the mission many between meaning of the mission of the mission many between meaning of the mission of the mission of the mission meaning of the mission of the mission of the mission meaning of the mission of the mission of the mission of the many of the mission of the mission of the many of the mission of the mission of the mission of the many of the mission of the mission of the mission of the mission of the mission of the mission	sample statistics, paired sample statistics or one-sample					
<ul> <li>hot test: analyze fixed-effects and random-effects measure, group description statistics, more sequences and the sequences with test of the same of pages processing and the sequences with test of the same of the same sequences and the se</li></ul>	test.					
<ul> <li>hot test: analyze fixed-effects and random-effects measure, group description statistics, more sequences and the sequences with test of the same of pages processing and the sequences with test of the same of the same sequences and the se</li></ul>	· ANOVA and ANCOVA: Conduct contrast, range, and post					
measure, group decripter, statistics, choose you model based on four hyse, of the suin of square procedure, parform licks of here thats, choose would be approximately consistant. Test to brainstee particular constants, or consistant. Test to brainstee particular constants, or the square same same. More partneric tests: Choose and decrip forges statistics of here thats, choose and decrip forges statistics. Confidence intervals for means, M. estimaters, interflucture of means, brainstee and decrip forges that the statistics. The same statistics requires two relations patients of here the statistics. Confidence intervals for means, M. estimaters, interflucture of means, brainstee and decrip forges that the statistics. The statistics in the statistics are discussed as a statistics. The statistics in the statistics of the statistics of the statistics of the statistics. The statistics in the statistics of the statistics in the scued as a statistics for that and there is remarker to any statistics for the statistics of the statistics. Statistics in the statistics of the statistics in the scued as a statistics for the statistics in the statistics and an algorithm that statistics with each case in a squarest chastistic based on statistics in the statistics and as algorithm that is statistics in the statistics makes as generated by the Provintiles procedure. Statistics and chastistics in the statistics in the statistics in the statistics in the statistics in the statistics in the statistics in the statistics in the statistics in the statistics in the statistics in the statistics in the statistics						
based on four types of the sum of squares proceedure perform lick of definition of adapters of proceedure performs. Act of definition of adapters of the proceedure sample, the independent samples, hindependent samples, the independent samples, hindependent periodes, high degree of floatishies, definition periodes, high degree of floatishies, definition periodes, high degree of floatishies, definition and new samples that the independent periodes and advecterities, using an algorithm that the independent periodes and advecterities, using an algorithm that the independent periodes, but which equals that the independent periodes and the independent samples. The independent periodes and the independent samples, high degree of floatishies and the independent samples of the independent periodes and the independent samples, high degree of floatishies and the independent samples and periodes that the independent samples and periodes that the independent and the independent samples. The independent samples and the independent samples, high degree of floatishies and the independent samples and periodes that the independent samples and the independent shares and periodes that the independent independent samples and the independent samples						
preferm tak of free treaty: choose balanced or insbalanced design: and analyse continues with public to mentods. Carnelistics: The first bioanize or partial carrelation, with researce. Nonparentic tests: Chi-square, Bioomial, Ruis, ees- sample, too indexe samples, residued samples, inclusioned isolation of the samples of the samples inclusioned isolation of the samples of the samples of the samples isolation of the samples of the samples of the samples isolation of the samples of the samples of the samples isolation of the samples of the samples of the samples isolation of the samples of the samples of the samples isolation of the samples of the samples of the samples isolation of the samples of the samples of the samples isolation of the samples of the samples of the samples isolation of the samples of the samples of the samples isolation of the samples of the samples of the samples isolation of the samples of the sample of the samples isolation of the samples of the sample of the sample of the samples isolation of the samples of the sample of the sample of the sample isolation of the samples of the sample of the sample of the sample of the sample isolation of the sample o						
design: and analyze covariance with up to 30 methods. Controllation: End to binaine a print location, or for distances. With programmer in the binaine a print location, and the sample, two independent samples, k-independent sample, two independent samples, k-independent samples, two independent samples, k-independent safe of location, interpaint potions safe of location, interpaint potions are samples and the print samples, k-independent safe of location, interpaint potions - There entities for first samples, k-independent safe of location, interpaint potions - There entities for interpaint, interpaint safe of location, interpaint safe of location, interpaint potions - There entities for interpaint, interpaint safe of location, interpaint potions - There entities of location, interpaint potions - Interpaint of location, interpaint potions - In						
Correlation: Test for bivariate or partial correlation, ruler distances indications is similarly of diseases is margine; two indexes disregations is the situations; description of orders; benefation of disregations of the situation of the situation of orders; benefations; the situations; description of orders; benefation; the situation; description of orders; benefation; the situation; description of orders; benefation; the situation; description of orders; benefation; the situation; and the situation of orders; benefation; and the situation; the situation; description; des						
distances indicating similarity or dissimilarity between measures. Honsamentric tests: Chisquare, binomial, Runs, ene- sample, too Independent samples, Nadeu denter tespice Confidences intervals for means: Mestimators: isterification of outcomes and identify forougs Fractor Analysic. Under of identify an unhead, or factors, that explain the pattern of contestion within a promak for ano orthogoal foronge intervalses, or factors, that explain the pattern of contestion within a promak for non-orthogoal foronge intervalses, rescent products. Subject of Non-orthogoal foronge intervalses. Subject of Non-orthogoal foronge intervalses. Subject on of two methods for namber of casses. Subject on of two methods for restrictly or classing an algorithm that can handle largen number of casses. Subject on orthogoal continue classes restrictly or classing an algorithm that starts with each test an asparate oblast and continue classes restrictly or classes. Subject on orthogoal classes restrictly classes. Subject on orthogoal classes restrictly or classes. Subject on orthogoal classes restrictly or classes. Subject on orthogoal classes restrictly or classes. Subject on orthogoal classes restrictly classes. The production classes restrictly classes. The production classes restrictly classes. The restrictly classes restrictly classes. The restrictl	design; and analyze covariance with up to 10 methods.					
measures. Horoparametric tests: Chi-square, Biocnial, Runs, one- sangle, two indesendent samples, k-independent samples, two indesendent samples, k-independent samples, two indesendent samples, k-independent samples, two index-samples, indexide samples. Hermitiation of cubies: Dotting of Indings. Tests to Predict. Numerical Outcome and Identify Groups factor Atalysis: Used to identify the underking variables, or factor Atalysis index to its indexide samples. Tests to Predict. Numerical Outcome and Identify Groups factor Atalysis: Used to identify transformer tests to Predict. Numbels for Inter analysis proceedure provana for non-Outbook for factors and provide Tests to Predict. Numbels for Inter analysis in the sweed Samples in Factor Samples. Tests to Predict. Numbels for Inter analysis in the sweed Samples for Inter analysis in the inter in the sweet Samples for Inter analysis in the inter i	·Correlation: Test for bivariate or partial correlation, or for					
measures. Horoparametric tests: Chi-square, Biocnial, Runs, one- sangle, two indesendent samples, k-independent samples, two indesendent samples, k-independent samples, two indesendent samples, k-independent samples, two index-samples, indexide samples. Hermitiation of cubies: Dotting of Indings. Tests to Predict. Numerical Outcome and Identify Groups factor Atalysis: Used to identify the underking variables, or factor Atalysis index to its indexide samples. Tests to Predict. Numerical Outcome and Identify Groups factor Atalysis: Used to identify transformer tests to Predict. Numbels for Inter analysis proceedure provana for non-Outbook for factors and provide Tests to Predict. Numbels for Inter analysis in the sweed Samples in Factor Samples. Tests to Predict. Numbels for Inter analysis in the sweed Samples for Inter analysis in the inter in the sweet Samples for Inter analysis in the inter i	distances indicating similarity or dissimilarity between					
<ul> <li>Idenzamentic tests: Chi-spare, Binomia, Rum, one-sample, too independent samples, too independent samples, too independent samples, Noneinsen Hervis too measures. Netwittators, Chi-spare Collidence Mercellis, Noneinsen Mercellis, N</li></ul>	· · · ·					
sample, two independent samples, k-independent samples, two independent samples, k-independent samples, two independent samples, k-independent samples, two index samples, k-independent samples, k-independent index samples, k-index samples, k-independent samples, k-independent index samples, k-independent index samples, k-index samples, k-independent samples, k-index samples, k-index samples, k-independent samples, k-index samples, k-index samples, k-independent samples, k-index samples, k-index samples, k-index samples, k-index samples, k-index samples, k-index samples, k-index samples, k-index samples, k-ind						
samples, two related samples, krelited samples. - Explore: Confidence introvis for means, Mestimator; tiertification of unlies; platting of findings. Exists to Predic Neuroidal Sacconses and denity Groups or factors, that explain the pattern of correlation within a set of observed variables. The factor analysis procedure provides a high degree of flexibility, offeng: - Seen methods of rotation, including direct oblimin and provides a high degree of flexibility, offeng: - an be saved as variables for further analysis. - an be saved as variables for further analysis. - and the saved as variables for tables of the saved as variables of noise - animer of classes. Select on os of totation, relatively homogenous groups of cases low which do for - relatively relatively relatively relatively relatively homogenous groups of cases low variables of choose for animity relatively relatively classifying only relatively relatively relatively classifying only relatively relatively relatively relatively relatively homogenous groups of cases low relative and combine clusters with each case in a separate cluster and combine clusters of an influence in the saved as variables of choose for animity relatively relatively and the saved as saved to help - you select the best solution. - Two Step Classes are displayed at each steps to help - you select the best solution. - How relatively relatively relatively relatively relatively - displayed at each step and/or in final surmary - output is displayed at each step and/or in final surmary - output is displayed at each step and/or in final surmary - ou						
<ul> <li>Lippiore: Confidence Interious for means, Mestimators; Identification of outless; potting of indings.</li> <li>Tasts to Predict: Numerical Jostcomes and Identify Groups Factor Analysis: Used to Identify the multiple of rotation, that explain the pattern of correlation within a promace in ship Appene of Installing, increase instance and the providence of the control of the correlation within a promace or non-orthogonal Totations.</li> <li>There embeds of computing factor scores. Also, scores can be saved as versibles for further analysis.</li> <li>Horman Call and the computing factor scores and the other instead disastriptic cancel and the other instead promace of the providence of the other instead disastriptic cancel.</li> <li>Hierarchield Call cancel and the other instead disastriptic cancel and disastriptic cancel and the other disastriptic cancel and disastriptic cancel and disastriptic cancel and the other disastriptic cancel and disastriptic cancel and the other disastriptic cancel and disastriptic and disastriptic cancel and disastriptic cancel and disastriptic and disastriptic cancel and disastrip</li></ul>						
idemitication of cultiers; plotting of findings. Factor: Analysis: Used to identify the underlying variables, or factors. That explain the pactre or correlation within a set of doeswell variables. The factor analysis procedure provides a high gene of finebility. - Five methods of computing factors cores. Also, scores - There methods of computing analysis. - There methods of computing analysis. - There methods of computing a algorithm that can handle large mometer of cases (or watch requires on the specific to which requires the specific to an algorithm that can handle large mometer of cases (or watch watch computed bases until only one is left. Analysis: Used to identify relatively more advices to subdation (gravofinations). Bistance procedures. Subtation: - Two Step Cluster Analysis: Group observations into clusters of left manufactor (arconsplate). - Two Step Cluster Analysis: Group observations to clusters of left manufactor (arconsplate). - There are constructs in the does adviced to restation. - There are person PLUM. Choose from seven options to clusters of left manufactors (marked methods in hand form. - Hierarchithe listication and R2 charge/peter di- specific the match the distater or theory in the specific to advice to watch the specific to advice the specific the distater methods: handward emination, for case, and cort of						
Tests Dredict: Numerical Outcomes and letently Groups - factor Analysis: Under to identify the underlying variables, or factors, that ocelian the pattern of correlation within as stor of based variables. The factors ranging proceedure process analysis degree of insubility direct oblinin and promax for non-orthogonal rations correlation within as promax for non-orthogonal rations correlations of the store						
<ul> <li>Factor Analysis: Used to identify the underlying variables, or factors, the degree of Hoobity, of ferring:</li> <li>Seen methods of factor extraction</li> <li>There methods of comparing factor scores. Also, scores can be aved as variables for there analysis.</li> <li>There methods of comparing factor scores. Also, scores can be aved as variables for there analysis.</li> <li>There methods of comparing factor scores. Also, scores can be aved as variables for there analysis.</li> <li>There methods of comparing factor scores. Also, scores can be aved as variables for there analysis.</li> <li>There methods of the which requires you to specify the number of classis which requires you to specify the number of classis which requires you to specify the number of classis of the variables for there analysis.</li> <li>Herarchical Clatter Analysis: Used to identify relatively nongenous groups of cases. For variables diverses until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and there is a specific baracteristic, using an algorithm that stars which excerts combines on the specific baracteristic, using an algorithm that stars and there analysis.</li> <li>Two Step Clutter Analysis: Group observations intro diverses and the number do closes to long.</li> <li>Two Step Clutter Analysis: Group observations intro diverses and the number do closes to long.</li> <li>Two Step Clutter Analysis: Group observations intro diverses and there analysis.</li> <li>Discrimitant: Offera a close of the variable sateletion of the specified produces numerous, long and entry.</li> <li>Continued Classes of the number do closes from a variable is distrified.</li> <li>Observation and k2 c</li></ul>	identification of outliers; plotting of findings.					
<ul> <li>Factor Analysis: Used to identify the underlying variables, or factors, the degree of Hoobity, of ferring:</li> <li>Seen methods of factor extraction</li> <li>There methods of comparing factor scores. Also, scores can be aved as variables for there analysis.</li> <li>There methods of comparing factor scores. Also, scores can be aved as variables for there analysis.</li> <li>There methods of comparing factor scores. Also, scores can be aved as variables for there analysis.</li> <li>There methods of comparing factor scores. Also, scores can be aved as variables for there analysis.</li> <li>There methods of the which requires you to specify the number of classis which requires you to specify the number of classis which requires you to specify the number of classis of the variables for there analysis.</li> <li>Herarchical Clatter Analysis: Used to identify relatively nongenous groups of cases. For variables diverses until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and combine classes until only one is left. Analysis: and there is a specific baracteristic, using an algorithm that stars which excerts combines on the specific baracteristic, using an algorithm that stars and there analysis.</li> <li>Two Step Clutter Analysis: Group observations intro diverses and the number do closes to long.</li> <li>Two Step Clutter Analysis: Group observations intro diverses and the number do closes to long.</li> <li>Two Step Clutter Analysis: Group observations intro diverses and there analysis.</li> <li>Discrimitant: Offera a close of the variable sateletion of the specified produces numerous, long and entry.</li> <li>Continued Classes of the number do closes from a variable is distrified.</li> <li>Observation and k2 c</li></ul>	Tests to Predict Numerical Outcomes and Identify Groups					
or factors, that explain the pattern of correlation within a set of observed analysis proceeding provides a high degree of floationity, offering: - Seen enables of factor extraction - File methods of factor extraction - File methods of factor extraction - Research enables is used to service the indigent of tables of sectors - Research enables is used to service the indigent of tables of sectors - Research enables is used to isenify relatively homogeneous groups of cases based on selected diardeer relatives you to specify the number of clusters. Select nee of two methods for disarching, cases, either updating cluster enters: intraview of classify and on the relative of the selected diardeer relative, workshipg only Hierarchial Cluster Analysis: Used to leader of the selected diardeer relatives and classify early with early of selection diardeer relatives of classify adjusted the there in the selection of the selection of the selection, with either diardeer relatives of classify adjusted the there of the selection of the selec						
set of observed virbables. The factor analysis procedure provides a high degree of factorize stration - Seven methods of factor extraction - Free methods of factor extraction - Three methods of factor extraction - Three methods of computing factor scores. Also, scores can be awed as viriables for three analysis. - Chronass Cluster Analysis: Used to identify relatively characterization, using an algorithm that can handle large number of clusters, Select one of two methods for disatifying cases, either updating cluster centers iteratively or classifying only. - Hierarchical Cluster Analysis: Used to identify relatively homogenous group of cases (or variables) based on selected characteristics, using an algorithm that starts with each case in a separate cluster and combine clusters unit only one is the Analysis: Used to identify relatively homogenous group of cases (or variables) based on selected character Analysis: Used to submit distarts with each case in a separate cluster and combine clusters unit only one is the Analysis are update backs based on a variety of standarding transformations. Distance group clusters are the number be chosen automatically. - Discriminant: Offers a choice of variables is backs of distarts or let the number be chosen automatically. - Discriminant: Offers a choice of variable selection method, statistic at each step and in final summary, output of displayed a cosh step and for final summary, output of displayed a cosh step and for fails summary, output of displayed a cosh step and for estimation, to specify numerical tolerance for checking singularly, and customes during the list factor case to used to specify the model. - Chronital Bereation For the scale on automatically. - Discriminant: Official regreesion-FUM: Choose from seven options to customic equity, free list factores and the relatively specified; paperty the blaster entry the lister of relation (with specified; paperty the blaster entry the lister specified; paperty the blaster entry the lister entry the lister specified;						
provides a high degree of feability, offering: - Seven methods of factor extraction - Five methods of contogonal rotations - Three methods of contogonal rotations - Three methods of computing factor scores. Also, scores can be aved as variables for futures ranyles. Homeson contegonal rotations - Three methods of computing factor scores. Also, scores can be aved as variables for futures ranyles. Homeson contegonal rotations - Three methods of computing hattor scores. Also, scores can be aved as variables for futures ranyles. Homeson contegonal rotations - Three methods of computing a lattor scores. Also, scores - mumber of cluster, Select on of two methods for clustifying cases, either updating cluster centers iteratively or classifying only. - Hierarchical Cluster Analysis: Used to identify relatively homogenous group of cases (or two methods for cluster future) of standarding transformations. Distance from a variety of standarding transformations. Distance form a variety of standarding transformations. Distance form a variety of standarding transformations. Distance form in the first a chuster analysis. Two Step Cluster Analysis. Group observations into clusters based to an earnees criterion, with ether categorial or continuous level data: specify the number of clusters in the cluster and in final summary: output is displayed at each step and in final summary: outputs displayed at each step and in final summary: outputs displayed at each step and in final summary: outputs displayed teach step and in final summary: outputs displayed teach step and in final summary: outputs displayed teach step and (or the mission to customis outputs. Intheran						
<ul> <li>Seven methods of factor extraction</li> <li>Free methods of rotation, including direct bilinin and proma for non-orthogonal rotations</li> <li>Three methods of computing factor scores. Also, scores can be saved as variables for further analysis.</li> <li>Kimeano Cluster Analysis: Used to identify relatively homogeneous groups of cases based on selected duration of the state of the selection of the sele</li></ul>	, ,					
- Five methods of rotation, including direct oblimin and promark for no-rotational controlling latter scores. Also, scores can be saved as variables for further analysis. Used to identify relatively homogeneous groups of cases based on selected characteristics, using an algorithm that can handle large number of clusters, Salect one of two methods for the number of clusters, Salect one of two methods for the number of clusters, Salect one of two methods for the number of clusters, Salect one of two methods for the number of clusters, Salect one of two methods for the number of cluster, Salect one of two methods for the number of cluster, Salect one of two methods for the number of cluster, Salect one of two methods for the number of cluster, Salect one of two methods for the number of cluster, Salect one of two methods for the number of cluster, Salect one of two methods for the number of cluster set of the number of cluster analysis. Used to identify relatively homogenous groups of cases (for variable) based on selected dharacteristic, using an algorithm that starts with each case in a separate cluster and combine clusters until only one is left. Analyze raw variables or choose for an avairely of task forup observations into clusters based on nearness circleno, with either case of allowing and task stage to help you select the best solution.  Two Step Cluster Analysis of Cluster Analysis, and Cluster, Salecter of significance; produces a numerous descriptive and equation statistic.  Ordinal regression-FLUK- Choose form seven options to control the transfer algorithm is clusters algorithm to clusters algorithm to clusters algorithm is cluster algorithm analy cluster set of the solutions, the set of the cluster algorithm and the clusters algorithm is cluster algorithm clusters and for the statistic.  Ordinal regression-FLUK- Choose form seven options to control the transfer algorithm used for estimation, to specify numerical tolerance for checking singlight, and to custome cluster for checking singlight, and to						
promator non-orthogonal rotations - Three methods of computing factors cores. Also, scores can be saved as variables for further analysis. - K-means Cluster Analysis: Used to identify relatively homogeneous groups of cases to a selected distancteristics, using an algorithm that can handle large number of cluster. Select on editors in that can handle large number of cluster. Select on editors in that can handle large number of cluster. Analysis: Used to identify relatively homogeneous groups of cases (or two methods for classifying cluster. Analysis: Used to identify relatively homogeneous groups of case (or two methods for classifying cluster. Analysis: Used to identify relatively homogeneous groups of case (or two methods for classifying cluster. Analysis: Used to identify relatively homogeneous groups of case (or two methods. Softmeter from a variety of standording transformations. Distance from a targeneous classifier and from. - Linear Regression. Choose from since naturatically. - Obscriminant. Offers a choice of variable selection relatives a set the number do from submatically. - Obscriminant. Offers a choice of variable selection official regression-FLUME. Choose from seven options to control the interview algorithm used for stimulation, to specific variables are traved as valid values. Distributes and for the selection (with a outcome specified), specify the distance metric used to measure specifical, specify the distance metric used to measure provements prependel Paper License with Authorization Code 12. morth MS Subscrytin	<ul> <li>Seven methods of factor extraction</li> </ul>					
<ul> <li>There methods of computing factor scores, Aloo, scores can be saved as variables for further analysis.</li> <li>"K-means Cluster Analysis: Used to identify relatively homogeneous groups of cases based on selected characteristics, using an algorithm that can handle large number of clusters, Select one of two methods for dashfying cases, either update you to specify the number of clusters, solect one diverse centers interactively or classifying only.</li> <li>Herarchical Cluster, Analysis: Used to identify relatively homogeneous groups of cases (or variables) based on selected characteristic, using an algorithm that starts with each case in a separate cluster and combine clusters until only one is fold. Analyzer any variables or choose from a variety of standardizing transformations. Distance or similarly measures are generated by the Proxomities procedure. Statistics are displayed at each stage to help you select the baracteristic, using an algorithm that into the unamber de characteristic, with and in final summary output: for the number of new automatically.</li> <li>Obsciminant: Offers a choice of variable selection method, tastistics at each step and in final summary output is displayed at each step and infinal summary output is displayed at each step and/or the theorem demony. forward entry, forward step wide selection and the change/set of selection and trace to the number of clusters and in final summary.</li> <li>Ordinal regression PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify the mixed selection (with segnificance: produces numerous descriptive and to such specific lay case) of cases; and control wether mixing values or categorical variable are treated as valid values.</li> <li>Dehember Derive Support (Sa): Fee urgrade for 1 year with technical support through a CSB.</li> <li>Diractical The provisioning of Technical Support Sa): Technical Support Sa): Technical Support Sa): Technical Support Sa): Technical Support</li></ul>	- Five methods of rotation, including direct oblimin and					
<ul> <li>There methods of computing factor scores, Aloo, scores can be saved as variables for further analysis.</li> <li>"K-means Cluster Analysis: Used to identify relatively homogeneous groups of cases based on selected characteristics, using an algorithm that can handle large number of clusters, Select one of two methods for dashfying cases, either update you to specify the number of clusters, solect one diverse centers interactively or classifying only.</li> <li>Herarchical Cluster, Analysis: Used to identify relatively homogeneous groups of cases (or variables) based on selected characteristic, using an algorithm that starts with each case in a separate cluster and combine clusters until only one is fold. Analyzer any variables or choose from a variety of standardizing transformations. Distance or similarly measures are generated by the Proxomities procedure. Statistics are displayed at each stage to help you select the baracteristic, using an algorithm that into the unamber de characteristic, with and in final summary output: for the number of new automatically.</li> <li>Obsciminant: Offers a choice of variable selection method, tastistics at each step and in final summary output is displayed at each step and infinal summary output is displayed at each step and/or the theorem demony. forward entry, forward step wide selection and the change/set of selection and trace to the number of clusters and in final summary.</li> <li>Ordinal regression PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify the mixed selection (with segnificance: produces numerous descriptive and to such specific lay case) of cases; and control wether mixing values or categorical variable are treated as valid values.</li> <li>Dehember Derive Support (Sa): Fee urgrade for 1 year with technical support through a CSB.</li> <li>Diractical The provisioning of Technical Support Sa): Technical Support Sa): Technical Support Sa): Technical Support Sa): Technical Support</li></ul>	promax for non-orthogonal rotations					
<ul> <li>can be saved as variables for further analysis.</li> <li>-K-means Cluster Analysis: Used to identify relatively</li> <li>homogeneous groups of cases based on selected</li> <li>draatceristics, using an algorithm that can handle large</li> <li>number of clusters, Select on of two methods for</li> <li>dassifying cases, either updating cluster centers</li> <li>iteratively or classifying only.</li> <li>- Hierarchical Cluster Analysis: Used to identify relatively</li> <li>homogenous groups of cases (or variables) based on</li> <li>selected characteristics, using an algorithm that starts</li> <li>with each case in a separate cluster and combine cluster</li> <li>until only one is left, Analyse raw variables or choose</li> <li>rowing starts are generated by the froimitties</li> <li>provedure. Statistics are displayed at each stage to help</li> <li>you select the best olution.</li> <li>Two step Cluster Analysis: Group observations into</li> <li>clusters based on nearness: choice of variable selection</li> <li>rethody, statistics are choice of variable selection</li> <li>rethody, statistics are choice of variable selection</li> <li>rethody, statistics are choice of variable selection</li> <li>rethody, statistics.</li> <li>Ordinal regression-PLUK. Choose from seven options to</li> <li>control the travel adjorith maximum, to</li> <li>respective heights.</li> <li>respective heights.</li> <li>respective heights and in fails summary;</li> <li>output displayed at each step and/or in fail form.</li> <li>-Ubscrimter adjority, the for prediction (with output effect of significance; produces numerous descriptive and equation statistics.</li> <li>Ordinal regression-PLUK. Choose from seven options to</li> <li>control the travel adjorithm used for estimation, to</li> <li>specify numerical tolerance for checking singularity, and</li> <li>to costoma coupt, the link functions can be used to</li> <li>specify numerical tolerance for checking singularity.</li> <li>Herarchield, specify the distance metric used to messure the similar</li></ul>	- Three methods of computing factor scores. Also, scores					
<ul> <li>-k.means Cluster Analysis: Used to identify relatively homogeneous groups of cases based on selected characteristics, using an algorithm that can handle large number of cases, either updating cluster centers iteratively or classifying group.</li> <li>-Hierarchical Cluster Analysis: Used to identify relatively homogeneous groups of cases (or variables) based on selected characteristics, using an algorithm that starts with each case in a separate cluster and combine clusters until only one is felf. Analyzer any variables or choose from a variety of standardizing transformations. Distance or similarly measures are generated by the Probamities procedure. Statistics are displayed at each stage to help you select the baracteristic, using an algorithm that entry clusters of aton meanses clusters.</li> <li>Two Step Cluster Analysis: Group observations into clusters of the number de charage and in final summary, output 5 displayed at each stage to help you select the baracteristic, using an algorithm that form clusters of the number de charage and in final summary, output 5 displayed at each stage and inform clusters and on meanses cluster, and in final summary, output 5 displayed at each stage and inform clusters and entry forcer termoval, finanzet entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and coution talteristics.</li> <li>Ordinal regression-PLW: Choose from seven options to control the iterative algorithm used for estimation, to specify unerical loterance for texting singularity, and to castorize output; five link functions can be used to specify unerical loterance of took (WAI); see upgrade for 1 year with technical support through a CSH.</li> <li>Pherentel Deparked Free traveling singularity, and to castorize output; five link functions can be used to specify unerical loterance of the clusting singularity, and to castorize output; five link functions can be used to specify the distance metric used to measure the similatify of casse; and control whet</li></ul>						
<ul> <li>homogeneous groups of cases based on selected characteristics, using an algorithm that can handle large number of cluster, Select on of two methods for classifying cases, either updating cluster centers iteratively or classifying const.</li> <li>Hierarchical Cluster Analysis: Used to identify relatively homogeneous groups of cases (or variables) based on selected characteristics, using an algorithm that starts with each case in a separate cluster and combine clusters in a variables or choose from a variety of standarding intradomations. Distance or similarity measures are generated by the Proximities procedure. Statistics are displayed at each stage to help you select the best solution.</li> <li>Two Step Cluster Analysis: Group observations into clusters output is displayed at each stage to help you select the best solution.</li> <li>Two Step Cluster Analysis: Group observations into clusters output is displayed at each stage not help you selection and relative and infini summary; outputs is displayed at each stage and in final summary; outputs is displayed at each stage and infinal summary; outputs is displayed at each stage and infini summary; outputs displayed at each stage notions be evidently.</li> <li>Cordinal regression-PLUM: Choose from six methods: backward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics.</li> <li>Ordinal regression-PLUM: Choose from six methods: backward entry, forward stepwise selection with each case and to nutcome specified; specify the elistance metric used to measure the similarity of cases; and control whether mising values or categorical variables are treated as valid values.</li> <li>Delevation Statistics.</li> <li>Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for a prediction (with specify outcome) or for classification (with outcome specified); specify the elistance metric used to measure the similarity of cases; and contol whether mising values o</li></ul>						
characteristics, using an algorithm that can handle large number of clusters, Select one of two methods for disasfing cases, either updating cluster centers iteratively or classifying govy. - Hierarchical Cluster Analysis: Used to identify relatively homogenous groups of cases (or variables) based on selected characteristics, using an algorithm that starts with each case in a separate cluster and combine clusters until only one is flef. Analyzer and variables or choose from a variety of standarding transformations. Distance or similarity measures are generated by the Proximities procedure. Statistics are displayed at each stage to help you select the based on nearness citreion, with either categorical or continuous level data, specify the number of clusters or best solution. - Two Step Cluster Analysis: Group observations into clusters based on nearness citreion, with either categorical or continuous level data, specify the number of clusters or best solution. - Uncar Regression: Choose from six methods: backward elimination, forget enroyal, forward entry, forward stepwide selection and a flag lummary; output is displayed at each step and/or in final form. - Linear Regression: Choose from six methods: backward elimination, forgories on lumores descriptive and equation statistics. - Ordinal regression: PLUM: Choose from seven options to control the iterative algorithm used for selimition, to specify hemotel. - Nearest Neighbor analysis: Used for prediction (with specify numotel, lostance netric used to measure specify bacterial lostrane for choose houses the subid values. - Disearded Bacteria support through a CSR - Discladed - Technical The provisioning of Technical Support - Support shall cover, at the minimum, the						
number of class but which requires you to specify the number of clusters. Select one of two methods for dassifying case, either updating cluster centers iteratively of classifying only. - Hierarchical Cluster Analysis: Used to identify relatively homogeneus groups of cases (or variable) so choose from a variety of standarding methods for clusters until only one is left. Analyze raw variables or choose from a variety of standarding methods on clusters until only one is left. Analyze raw variables or choose from a variety of standarding methods. Distance or similarity measures are generated by the Proximities procedure. Statistics are displayed at each stage to help you select the best solution. - Two Step Cluster Analysis: Group observations into clusters or let the number be chosen automatically. - Discriminant: Offers a choice of variable selection method, statistics at each step and in final summary; output is displayed at each step and/or in final form. - Linear Regression: Choose from six methods: backward elimination, fore denty, forced ennoy, forward enty, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. - Ordinal regression-PLUM: Choose from sixen options to control the iterative algorithm used for estimation, to specify humerical lobrance for checking singularity, and to customize output; five link functions can be used to specify the odistance metric used to measure the similarity of cases; and control whether mising values or categorical variables are treated as valid values. - Delavaelise Preptual Paper License with - Authorization Code for Liyaen with technical support through a CSR - CD included						
number of clusters, Select one of two methods for classifying cases, either updating cluster centers iteratively or classifying only. - Hierarchical Cluster Analysis: Used to identify relatively homogenous groups of cases (or variables) based on selected characteristics, using an algorithm that starts with each case in a separate cluster and combine clusters until only one is left. Analyse are variables or choose from a variety of standarding transformations. Distance or similarity measures are generated by the Provinities procedure. Statistics are displayed at each stage to help vou select the base solution. 'Two Step Cluster Analysis: Group observations into clusters based on nearness criterion, with either categorical or continuous level data; specify the number of clusters or based on nearness criterion, with either categorical or continuous level data; specify the number of clusters or based on nearness criterion, with either categorical at each step and in final summary; output is displayed at each step and/or in final form. -Linear Regression: Choose from six methods: backward elimination, forede entry, forever enrowal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordnal regression-RLWI: Choose from seven options to control the iterative algorithm used for estimation, to specify the model. -Nearest Neighbor analysis: Used for prediction (with specify classe; and control whether missing values or categorical variables are treated a valid values. Deiverview produces numers with the outcate the save of specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated a valid values. Deiverview Proper License with Authorization Code 12-month WS valuerytorical Support (KSS); Free uggrade for 1 year with technical Support (KSS); Free uggrade for year with technical Support (KSS); Free uggrade for year with techni						
<ul> <li>classifying cases, either updating cluster centers</li> <li>literatively or classifying only.</li> <li>· Hierarchical Cluster Analysis: Used to identify relatively</li> <li>homogenous groups of cases (or variable) based on</li> <li>selected characteristics, using an algorithm that starts</li> <li>with each case in a separate cluster and combine clusters</li> <li>until only one is left. Analyze raw variables or choose</li> <li>from a varlety of standordilling transformations. Distance</li> <li>or similarity measures are generated by the Proximities</li> <li>procedure. Statistics are displayed at each stage to help</li> <li>vou select the basis clutton.</li> <li>Two Step: Cluster Analysis: Group observations into</li> <li>clusters based on nearness criterion, with either</li> <li>categorical or continuous level data; specify the number</li> <li>of clusters or lased on nearness criterion, with either</li> <li>categorical or continuous level data; specify the number</li> <li>of clusters or lase on the number be chosen automatically.</li> <li>Discriminant: Offers a choice of variable selection</li> <li>method, statistics at each step and/or in final form.</li> <li>-linear Regression: Choose from seven options to</li> <li>control the lentware bet clustor.</li> <li>Ordinal regression:PUW: Choose from seven options to</li> <li>control the lentware data previous data based to specify the model.</li> <li>values or categorical action whether missing</li> <li>values or categorical action whether missing</li> <li>values or categorical availables are treated as valid values.</li> <li>bainerskie, specifical support (Bas): Free upgrade</li> <li>for it, are with technical support (Bas): Free upgrade</li> <li>for it, are with technical support (Bas): Free upgrade</li> <li>for it, are with technical Support (Bas): Free upgrade</li> <li>for it, are with technical Support (Bas): Free upgrade</li> <li>for it, are with technical Support (Bas): Free upgrade</li> <li>for it, are with technical Support (Bas): Free upgrade</li></ul>	number of cases but which requires you to specify the					
<ul> <li>Iteratively or classifying only.</li> <li>Hierarchical Guster Analysis: Used to identify relatively homogenous groups of cases (or variables) based on selected characteristics, using an algorithm that starts with each case in a separate cluster and combine clusters until only one is left. Analyze raw variables or choose from a variety of standardizing transformations. Distance or similarity measures are generated by the Proximites procedure. Statistics are displayed at each stage to help you select the best solution.</li> <li>Two Step Cluster Analysis: Group observations into clusters based on mearness criterion, with either categorical or continuous level data, specify the number of clusters or let the number be chosen automatically. Discriminant: Offers a choice of variable selection method, statistics at each step and in final summary, output is displayed at each step and/or in final form.</li> <li>Linear Regression: Choose from six methods: backward eithin, forward stepwise selection and R2 change/test of significance; produces numerous discriptive and equation stitistics.</li> <li>Ordinal regression-PUML: Choose from seven options to control to level bits functions can be used to specify the model.</li> <li>Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the disclase are trated as valid values.</li> <li>Deliwarbias Perpetual Paper License with Authorization Code</li> <li>12-month SW subscription &amp; Support (S&amp;S): Free upgrade for 1 year with technical support through a CS.</li> <li>Dincluded</li> <li>Technical The provisioning of Technical Support</li> <li>Support shall cover, at the minimum, the</li> </ul>	number of clusters, Select one of two methods for					
<ul> <li>Iteratively or classifying only.</li> <li>Hierarchical Guster Analysis: Used to identify relatively homogenous groups of cases (or variables) based on selected characteristics, using an algorithm that starts with each case in a separate cluster and combine clusters until only one is left. Analyze raw variables or choose from a variety of standardizing transformations. Distance or similarity measures are generated by the Proximites procedure. Statistics are displayed at each stage to help you select the best solution.</li> <li>Two Step Cluster Analysis: Group observations into clusters based on mearness criterion, with either categorical or continuous level data, specify the number of clusters or let the number be chosen automatically. Discriminant: Offers a choice of variable selection method, statistics at each step and in final summary, output is displayed at each step and/or in final form.</li> <li>Linear Regression: Choose from six methods: backward eithin, forward stepwise selection and R2 change/test of significance; produces numerous discriptive and equation stitistics.</li> <li>Ordinal regression-PUML: Choose from seven options to control to level bits functions can be used to specify the model.</li> <li>Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the disclase are trated as valid values.</li> <li>Deliwarbias Perpetual Paper License with Authorization Code</li> <li>12-month SW subscription &amp; Support (S&amp;S): Free upgrade for 1 year with technical support through a CS.</li> <li>Dincluded</li> <li>Technical The provisioning of Technical Support</li> <li>Support shall cover, at the minimum, the</li> </ul>	classifying cases, either updating cluster centers					
Hierarchical Cluster Analysis: Used to identify relatively homogenous groups of cases (or variable) based on selected characteristics, using an algorithm that starts with each case in a separate cluster and combine clusters until only one is left. Analyze raw variables or choose from a variety of standarding transformations. Distance or similarity measures are generated by the Proximities procedure. Statistics are displayed at each stage to help you select the best solution. "Two Step Cluster Analysis: Group observations into clusters based on nearness criterion, with either categorical or continuous level data; specify the number of clusters or step and in final summary output is displayed at each step and in final summary output is displayed at each step and in final summary output is displayed at each step and/or in final form. - Linear Regression: Choose from six methods: Eackward elimination, forced entry, forced removal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. Ordinal regression: House from seven options to control the tractive algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specified; progetify analysis: Used for prediction (with specified) specify the distance metric used to measure the similarity of cases; and control whether missing volues or categorical variables are treated as valid values. Discuster Experimed is support (SAS); free upgrade for 1 year with technical support (SAS); free upgrade for 1 year with technical support through a CSR CD include Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
<pre>homogenous groups of cases (or variables) based on selected characteristics, using an algorithm that starts with each case in a separate cluster and combine clusters until only one is left. Analyze raw variables or choose from a variety of standarding transformations. Distance or similarity measures are generated by the Proximites procedure. Statistics are displayed at each stage to help you select the best solution. Two Step Cluster Analyzes: Group observations into clusters based on nearness criterion, with either categorical or continuous level dats; specify the number of clusters or let the number be chosen automatically. Discriminant: Offers a choise of variable selection method, statistics at each step and in final summary; output is displayed at each step and form. .tunear Regression: Choose from six methods: backward elimination. Cored entry, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistic. Ordinal regression+ULM: Choose from seven options to control the trative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; the link functions can be used to specify the model. Nearest Neighbor analysis: Used for prediction (with specified): specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription &amp; Support (S&amp;S): Free upgrade for 1 year with technical support through a CSN CD included Technical The provisioning of Technical Support Support Shall cover, at the minimum, the</pre>						
selected characteristics, using an algorithm that starts with each case in a separate cluster and combine clusters until only one is left. Analyze raw variables or choose from a variety of standardizing transformations. Distance or similarity measures are generated by the Proximities procedure. Statistics are displayed at each stage to help you select the best solution. "Two Step Cluster Analysis: Group observations into clusters bases criterion, with either categorical or continuous level data; specify the number of clusters or bases criterion, with either categorical or continuous level data; specify the number of clusters or bases criterion, with either categorical are chosen automatically. Obscriminant: Offers a choice of variable selection method, statistics at each step and/or in final form. -Linear Regression: Choose from six methods: backward elimination, forced entry, forcard entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify the model. Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specify the model. Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specify the singer treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (SaS): Free upgrade for 1 year with technical support through a CSR CD include Technical The provisioning of Technical Support						
with each case in a separate duster and combine clusters until only one is left. Analyzer aw variables or choose from a variety of standardizing transformations. Distance or similarity measures are generated by the Proximities procedure. Statitics are deshaped at each stage to help you select the best solution. Two Step Cluster Analysis: Group observations into clusters based on nearness criterion, with either categorical or continuous level data; specifi the number of clusters or let the number be chosen automatically. Observations and the specific data; specifi the number of clusters or let the number be chosen automatically. Output is displayed at each step and in final summary; output is displayed at each step and or final summary; output is displayed at each step and for infinal form. Unear Regression: Choose from six methods: backward elimination, forced entry, forced removal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the distance metric used to measure the similarity of classification (with specific outcome) or for classification (with specific outcome) or for classification (with specific dustistics) are treated as valid values. Deliverables Perpetual Paper License with Authorization Gode 12-month SW Subscription & Support (SSs): Free upgrade for 1 year with technical support through a CSR CD included						
until only one is leff. Analyze raw variables or choose from a variety of standarding transformations. Distance or similarity measures are generated by the Proximities procedure. Statistics are displayed at each stage to help you select the best solution. Two Step Cluster Analysis: Group observations into clusters based on nearness criterion, with either categorical or continuous level data; specify the number of clusters or bases criterion, with either categorical or continuous level data; specify the number of clusters or bases criterion, with either categorical or continuous level data; specify the number of clusters or bases criterion, with either categorical or continuous level data; specify the number of clusters or development and summary; output is displayed at each step and in final summary; output is displayed at each step and or in final form. -Linear Regression: Choose from six methods: backward elimination, forced entry, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking signily and to customize output; five link functions can be used to specify the model. -Narrest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specify be model. -Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (SAS): Free ungrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the	selected characteristics, using an algorithm that starts					
<pre>from avariety of standardizing transformations. Distance or similarity measures are generated by the Proximities procedure. Statistics are displayed at each stage to help you select the best solution. Two Step Cluster Analysis: Group observations into clusters based on nearness criterion, with either categorical or continuous level data; specify the number of clusters or let the number be chosen automatically. Discriminant: Offers a choice of variable selection method, statistics at each step and in final summary; output is displayed at each step and/or in final form. - uncer Regression: Choose from six methods: backward elimination, forced entry, forced removal, forward entry, forward stepsies selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the literative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the odel. -Nearest. Neighbor analysis: Used for prediction (with specific outputs are treated as valid values. Deliverables Perpletual Perpletual Paper License with Authorization Code 12-month SW Subscription &amp; Support (S&amp;S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the</pre>	with each case in a separate cluster and combine clusters					
or similarity measures are generated by the Proximities procedure. Statistics are displayed at each stage to help you select the best solution. Two Step Cluster Analysis: Group observations into clusters based on nearnes criterion, with either categorical or continuous level data, specify the number of clusters or let the number be chosen automatically. -Discriminant: Offers a choice of variable selection method, statistics at each step and/or in final summary; output is displayed at each step and/or in final form. -Linear Regression: Choose from six methods: backward elimination, forced entry, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specified); specify the distance metric used to measure the similarity of case; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with - Authorization Code 12-month SW Subscription & Support (S&S): Free ugrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the	until only one is left. Analyze raw variables or choose					
or similarity measures are generated by the Proximities procedure. Statistics are displayed at each stage to help you select the best solution. Two Step Cluster Analysis: Group observations into clusters based on nearnes criterion, with either categorical or continuous level data, specify the number of clusters or let the number be chosen automatically. -Discriminant: Offers a choice of variable selection method, statistics at each step and/or in final summary; output is displayed at each step and/or in final form. -Linear Regression: Choose from six methods: backward elimination, forced entry, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specified); specify the distance metric used to measure the similarity of case; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with - Authorization Code 12-month SW Subscription & Support (S&S): Free ugrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the	from a variety of standardizing transformations. Distance					
procedure. Statistics are displayed at each stage to help you select the best solution. Two Step Cluster Analysis: Group observations into clusters based on mearness criterion, with either categorical or continuous level date; specify the number of clusters or let the number be chosen automatically. Discriminant: Offers a choice of variable selection method, statistics at each step and/or in final summary; output is displayed at each step and/or in final form. -linear Regression: Choose from six methods: backward elimination, forced entry, forced removal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverable: Perptual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free ungrade for 1 year with technical support though a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
you select the best solution. Two Step Cluster Analysis: Group observations into Clusters based on nearness criterion, with either categorical or continuous level data; specify the number of clusters of let the number be chosen automatically. Discriminant: Offers a choice of variable selection method, statistics at each step and/or in final summary; output is displayed at each step and/or in final form. Linear Regression: Choose from six methods: backward elimination, forced entry, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. Ordinal regression-PLUM: Choose from seven options to control the trative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverable: Perptual Paper License with Authorization Code 12-month SW subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
<ul> <li>Two Step Cluster Analysis: Group observations into clusters based on nearness criterion, with either categorical or continuous level data; specify the number of clusters or let the number be chosen automatically.</li> <li>Discriminant: Offers a choice of variable selection method, statistics at each step and in final summary; output is displayed at each step and/or in final form.</li> <li>Linear Regression: Choose from six methods: backward elimination, forced entry, fored removal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics.</li> <li>Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model.</li> <li>Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values.</li> <li>Deliverables Perpetual Paper License with Authorization Code</li> <li>12-month SW Subscription &amp; Support (S&amp;S): Free upgrade for 1 year with technical support (S&amp;S): Free upgrade for 1 year with technical support (S&amp;S): Free upgrade for 1 year with technical support (S&amp;S): Free upgrade for 1 year with technical support through a CSR</li> <li>CD included</li> </ul>						
<pre>clusters based on nearness criterion, with either categorical or continuous level data; specify the number of clusters or let the number be chosen automatically. -Discriminant: Offers a choice of variable selection method, statistics at each step and/or in final summary; output is displayed at each step and/or in final form. -Linear Regression: Choose from six methods: backward elimination, forced entry, forced removal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. -Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specifie outcome) or for classification (with no outcome specifie outcome) or for classification (with no outcome specifie outcome) are treated as vaid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription &amp; Support (S&amp;S): Free upgrade for 1 year with technical Support (S&amp;S): Free upgrade for 1 year with technical Support (S&amp;S): Free upgrade for 1 year with technical Support fucung a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the</pre>						
categorical or continuous level data; specify the number of clusters or let the number be chosen automatically. Discriminant: Offers a choice of variable selection method, statistics at each step and/or in final form. -Linear Regression: Choose from six methods: backward elimination, forced entry, forced removal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. -Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specific outcome) or for classification (with no outcome specific outcome) or for classification (with measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support (hrough a CSR CD included Technical The provisioning of Technical Support						
of clusters or let the number be chosen automatically. ·Discriminant: Offers a choice of variable selection method, statistics at each step and in final summary; output is displayed at each step and/or in final form. -Linear Regression: Choose from six methods: backward elimination, forced entry, forced removal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. -Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specifix outcome) or for classification (with no outcome the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
<ul> <li>Discriminant: Offers a choice of variable selection method, statistics at each step and/or in final summary; output is displayed at each step and/or in final form.</li> <li>-Linear Regression: Choose from six methods: backward elimination, forced entry, forced removal, forward escipation and R2; change/test of significance; produces numerous descriptive and equation statistics.</li> <li>Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model.</li> <li>Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or classification ds Support Subscription &amp; Support (S&amp;S): Free upgrade for 1 year with technical support (S&amp;S): Free upgrade for 1 year with technical support mough a CSR</li> <li>CD included</li> <li>Technical The provisioning of Technical Support</li> </ul>						
method, statistics at each step and in final summary; output is displayed at each step and/or in final form. -Linear Regression: Choose from six methods: backward elimination, forced entry, forced removal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. -Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specifie output; five distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SS Subscription & Support (S&S): Free upgrade for 1 year with technical support (for schild Support Support shall cover, at the minimum, the	of clusters or let the number be chosen automatically.					
method, statistics at each step and in final summary; output is displayed at each step and/or in final form. -Linear Regression: Choose from six methods: backward elimination, forced entry, forced removal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. -Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specifie output; five distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SS Subscription & Support (S&S): Free upgrade for 1 year with technical support (for schild Support Support shall cover, at the minimum, the	·Discriminant: Offers a choice of variable selection					
output is displayed at each step and/or in final form.         -Linear Regression: Choose from six methods: backward         elimination, forced entry, forced removal, forward entry,         forward stepwise selection and R2 change/test of         significance;       produces numerous descriptive and         equation statistics.         Ordinal regression-PLUM: Choose from seven options to         control the iterative algorithm used for estimation, to         specify numerical tolerance for checking singularity, and         to customize output; five link functions can be used to         specify numerical tolerance metric used to measure         the similarity of cases; and control whether missing         values or categorical variables are treated as valid values.         Deliverables       Perpetual         Paper       License         Lizenonth SV Subscription & Support (S&S): Free upgrade         for 1 year with technical support through a CSR         CD included         Technical       The provisioning of Technical Support         Support       shall cover, at the minimum, the	method, statistics at each step and in final summarv:					
-Linear Regression: Choose from six methods: backward elimination, forced entry, forced removal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. -Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of crase; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
elimination, forced entry, forced removal, forward entry, forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify numerical tolerance for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support						
forward stepwise selection and R2 change/test of significance; produces numerous descriptive and equation statistics. Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specifi the model. Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
significance; produces numerous descriptive and equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. -Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
equation statistics. -Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. -Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
-Ordinal regression-PLUM: Choose from seven options to control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. -Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. -Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the	equation statistics.					
control the iterative algorithm used for estimation, to specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. -Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the	·Ordinal regression-PLUM: Choose from seven options to					
specify numerical tolerance for checking singularity, and to customize output; five link functions can be used to specify the model. •Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
to customize output; five link functions can be used to specify the model. •Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the	-					
specify the model. •Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
<ul> <li>Nearest Neighbor analysis: Used for prediction (with specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values.</li> <li>Deliverables Perpetual Paper License with Authorization Code</li> <li>12-month SW Subscription &amp; Support (S&amp;S): Free upgrade for 1 year with technical support through a CSR CD included</li> <li>Technical The provisioning of Technical Support Support shall cover, at the minimum, the</li> </ul>						
specific outcome) or for classification (with no outcome specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
specified); specify the distance metric used to measure the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
the similarity of cases; and control whether missing values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the	specified); specify the distance metric used to measure					
values or categorical variables are treated as valid values. Deliverables Perpetual Paper License with Authorization Code 12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the	the similarity of cases; and control whether missing					
Deliverables       Perpetual       Paper       License       with         Authorization Code						
Authorization Code         12-month SW Subscription & Support (S&S): Free upgrade         for 1 year with technical support through a CSR         CD included         Technical       The provisioning of Technical Support         Support       shall cover, at the minimum, the	÷					
12-month SW Subscription & Support (S&S): Free upgrade for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
for 1 year with technical support through a CSR CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
CD included Technical The provisioning of Technical Support Support shall cover, at the minimum, the						
Technical         The provisioning of Technical Support           Support         shall         cover, at the minimum, the						
Support shall cover, at the minimum, the						
Support shall cover, at the minimum, the	Technical The provisioning of Technical Support					
		1	I			

<ul> <li>Help with errors during installation</li> <li>Help with failed installation or when program does not run</li> <li>On-site installation of software</li> <li>Provide guidance in encryption codes</li> <li>Provide details on recommended resource settings and how to set them</li> <li>Walk through update of encryption code, providing PLM with step-by-step instructions over the phone</li> <li>In situations in which common troubleshooting techniques have been unsuccessful, initial step is to walkthrough the installation procedure with the customer; If necessary, the issue shall be escalated to an on-site troubleshooting by a CSR representative.</li> </ul>				
Training As a Change Management Program, training programs on the optimal use of the Mathematical Software shall be administered to at least two (2) PLM employees who shall render effective services to the University on the use thereof.				
	Grand Tota	al Price Php		

This is to certify that I personally conducted the canvass and that the data herein are true and correct.

## (Signature over Printed Name) Landline: Mobile No.: E-Mail:

### **RON MARR P. EVANGELISTA**

Canvasser

# **TERMS AND CONDITIONS:**

- 1. Bidders shall provide correct and accurate information required in this form.
- 2. Bidders may quote for any or all the items.
- 3. Price quotation/s must be valid for a period of *Thirty (30) calendar days* from the date of submission.
- 4. Price quotation/s, to be denominated in Philippine Peso shall include all taxes, duties and/or levies payable subject to deduction of applicable Government Tax. PLM is a Government Institution, as such it is mandated to withhold taxes and issue the corresponding BIR certification (BIR Form #2306, 2307).Computations and rates are as follows:

a. For VAT Registered

(PHP10,000.00 and below)

· · ·		,	
	FINAL VAT	EWT	TOTAL
GOODS	5%	0%	5%
SERVICES	5%	0%	5%

(Above PHP10,000.00)

	FINAL VAT	EWT	TOTAL
GOODS	5%	1%	6%
SERVICES	5%	2%	7%

b.	For NON-VAT Registered

(PHP10,000.00 and below)

<u> </u>	/		
	PERCENTAGE	EWT	TOTAL
GOODS	3%	0%	3%
SERVICES	3%	0%	3%

(Above PHP10,000.00)

	PERCENTAGE	EWT	TOTAL
GOODS	3%	1%	4%
SERVICES	3%	2%	5%

- 5. Quotations exceeding the Approved Budget for the Contract shall be rejected.
- 6. Award of the Contract shall be made to the lowest quotation (for goods and infrastructure) or, the highest rated offer (for consulting services) which complies with the minimum technical specifications and other terms and conditions stated herein.
- 7. Any interlineations, erasures or overwriting shall be valid only if they signed or initiated by you or any of your duly authorized representative/s.
- 8. The item/s shall be delivered within \_\_\_\_\_\_ working days from the receipt of the approved Purchase Order (PO).
- 9. PLM shall have the right to inspect and/or to test the goods to confirm their conformity to the technical specifications.
- 10. Liquidated damages equivalent to the one tenth of one percent (0.001%) of the value of the goods not delivered within the prescribed delivery period shall be imposed per day of delay. PLM shall rescind the contract once the cumulated amount of liquidated damages reached ten percent (10%) of the amount of the contract, without prejudice to other courses of action and remedies open to it.